



Water-Level Changes at Devil's Hole Associated with Barometric Pressure, Earth Tides and Earthquakes

June 2, 2004

Paula A. Cutillo, Ph.D.

Hydrogeologist

Water Resources Division, Water Rights Branch

National Park Service



Collaborators

- Keith Halford, USGS, Nevada District
- Randy Laczniaak, USGS, Nevada District



Outline


- Effects of atmospheric pressure and earth tides on water levels
- Filtered Devil's Hole water-level record
- Effects of earthquakes on water levels

National Park Service



EXPERIENCE
YOUR
AMERICA

Monthly Mean Water Level (Jun-62 to Mar-04)




National Park Service



Hourly & Daily Mean Water Level (Oct-89 to Mar-04)

EXPERIENCE
YOUR
AMERICA





Factors Influencing Water Level


- Atmospheric pressure changes
- Earth tides
- Crustal strain (e.g., earthquakes)
- Ground-water development
- Climate change

National Park Service



EXPERIENCE
YOUR
AMERICA

Barometric Pressure & Water Level during the 1999 Hector Mine Earthquake





Methods


- **Data requirements**
 - Hourly BP and WL time series
 - Theoretical Earth tide time series (Harrison, 1971)
- **Analyze how water level is affected by earth tides and changes in barometric pressure**
 - Used Visual Basic fitting program (developed by K. Halford, USGS) that estimates a **multiplier** that changes the amplitude and a **constant** that shifts the phase of each time series response
 - Used small data sets (15-30 day time periods)
 - Check for consistency over time




Methods

- **Filter entire data set**
 - Used average multiplier and constant
 - Developed a Fortran program to process over 10 years of hourly data
 - Generated water-level record filtered for the effects of barometric pressure and earth tides

Measured & Filtered Daily Mean Water Level




Surrogate & Filtered Daily Mean Water Level




Earth Tide Sensitivity

6-month period – hourly water level


**Filtered for barometric
pressure only**



Filtered for Earth tides only



Measured & Filtered Daily Water Level






Summary & Conclusions

- Water-level fluctuations due to air pressure and earth tides do not exceed 0.4 ft at Devil's Hole
- Installation of a barometer
- Filtered water level data helps to isolate other factors that influence water levels (e.g., earthquakes)

Earthquake epicenters






Earthquakes

- Common hydrogeologic responses
 - Water-level oscillations due to seismic waves (dynamic strain)
 - Instantaneous water-level offsets due to aquifer compression/expansion (volumetric strain)
- Factors influencing response
 - Size, depth, distance from epicenter



- Seismic Waves

- Cause water-level oscillations resembling seismograms
- Only detected when water levels are recorded at very small (e.g., 1 s) intervals




National Park Service




EXPERIENCE
YOUR
AMERICA

- **Crustal Strain**

- Fault movement causes rock compression and expansion
- Coseismic +/- offset or step change in water level



**1992
Landers/LSM:
 $\Delta WL > 0.7$ ft**



**1999
Hector Mine:
 $\Delta WL = 0.1$ ft**




Methods

- **Are the coseismic water-level offsets a result of the strain field produced by the earthquakes?**
 - Calculated volumetric strain field using known earthquake parameters and a dislocation model (Okada, 1992)
 - Converted change in strain to change in water level (Ge and Stover, 2000)

1992 Landers earthquake

$M_w=7.3$


Right-lateral strike-slip



1992 Little Skull Mountain earthquake

$$M_w=5.6$$


Normal fault



1999 Hector Mine earthquake

$M_w=7.0$

Right-lateral strike-slip







Summary & Conclusions

- Coseismic water-level offsets agree with modeled volumetric strain in sign and magnitude
- Installation of strain meter(s)
- Future work - understand postseismic recovery

Postseismic Recovery - Little Skull Mtn. Earthquake



Filtered daily mean
water level at
Devil's Hole



Simulated water
level near LSM fault
zone



Conclusions

- Water-level fluctuations due to air pressure and earth tides are relatively small
- Coseismic offsets are likely a result of volumetric strain
- Installation of barometer and strain meter(s) will aid in filtering future water levels, and will aid in understanding postseismic water-level recovery



Acknowledgements

- NPS, Water Rights Branch
 - Chris Gable, Brad Gillies
- NPS, Death Valley National Park
 - Terry Fisk, Jim Roche